

Tank Mixing Compatibilities

Please read the following before considering using any of the listed tank mixes with BotaniGard WP:

This Compatibility List provides a record of physical compatibility tests that have been carried out with the listed products in a 2-way tank mix with BotaniGard WP, unless otherwise stated. No testing of biological compatibility, crop safety or efficacy has been undertaken. Use is entirely at user's own risk and no warranty is given or liability accepted by Certis.

Since testing Certis cannot guarantee that every entry on the Compatibility List currently has a valid approval. The occurrence of a particular mix on the list does not, therefore, imply or endorse the legal status of that mix and product labels should always be consulted for each product in the first instance. Other label recommendations on the product labels should also be adhered to.

Ensure constant agitation from mixing to spraying, which should occur immediately after mixing is completed.

FUNGICIDES

<u>BRAND NAME</u>	<u>ACTIVE INGREDIENT</u>	<u>COMMENTARY</u>
Amistar®	Azoxystrobin	Observe a spray interval of 4 days before & after application
AQ10®	<i>Ampelomyces quisqualis</i> strain AQ10	Physically compatible
Bravo®	Chlorothalonil	Observe a spray interval of 4 days before & after application
Bumper® 250	Propiconazole	Observe a spray interval of 2 days before & after application
Captan® 80WG	Captan	Observe a spray interval of 4 days before & after application
Cercobin® WG	Thiophanate methyl	Physically compatible
Cuprolyt®	Copper oxychloride	Observe a spray interval of 4 days before & after application
Dithane® 945	Mancozeb	Observe a spray interval of 4 days before & after application
Fenomenal®	Fenamidone + fosetyl-aluminium	Physically compatible
Fytosave®	COS-OGA	Physically compatible



Flexity®	Metrafenone	Physically compatible
Folicur®	Tebuconazole	Observe a spray interval of 4 days before & after application
Frupica® SC	Mepanipyrim	Physically compatible
Gavel®	Imazalil	Observe a spray interval of 2 days before & after application
Karma®	Potassium hydrogen carbonate	Physically compatible
Luna Privilege®	Fluopyram	Physically compatible
Microthiol®	Sulfur 80%	Observe a spray interval of 2 days before & after application
Nimrod®	Bupirimate	Physically compatible
Paraat®	Dimethomorph	Physically compatible
Previcur Energy®	Propamocarb hydrochloride + fosetyl-aluminium	Physically compatible
Prolectus®	Fenpyrazamine	Physically compatible
Scala®	Pyrimethanil	Physically compatible
Serenade® ASO	<i>Bacillus subtilis</i> strain QST 713	Physically compatible
Signum®	Boscalid and pyraclostrobin	Observe a spray interval of 4 days before & after application
Stroby® WG	Kresoxim-methyl	Observe a spray interval of 2 days before & after application
Subdue®	Metalaxyl-M	Physically compatible
Sulfur volatilized	Sulphur	Observe a spray interval of 2 days before & 3 days after application
Sulfur - wet spray	Sulphur	Physically compatible: Observe a spray interval of 4 days before & after application



Switch®	Cyprodinil + Fludioxonil	Observe a spray interval of 4 days before & after application
Takumi®	Cyflufenamid	Observe a spray interval of 2 days before & after application
Teldor®	Fenhexamid	Physically compatible
Trianium P®	<i>Trichoderma harzianum</i> Rifai strain T-22	Physically compatible
Topas®	Penconazole	Observe a spray interval of 2 days before & after application
Topsin®	Thiophanate-methyl	Physically compatible

INSECTICIDES

<u>BRAND NAME</u>	<u>ACTIVE INGREDIENT</u>	<u>COMMENTARY</u>
Actara®	Thiamethoxam	Physically compatible
Applaud® 25SC	Buprofezin	Physically compatible Constant agitation advised
Azatin®	Azadirachtin-A	Physically compatible
Batavia®	Spirotetramat	Physically compatible
Dimilin®	Diflubenzuron	Physically compatible
Dipel®	<i>Bacillus thuringiensis</i> spp. <i>kurstaki</i>	Physically compatible
Dynamec®	Abamectin	Yes, but <u>spray immediately</u>
Kanemite®	Acequinocyl	Physically compatible
Majestik® / Eradicoat®	Maltodextrin	Physically compatible
Merit Forest®	Imidacloprid	Physically compatible
Prev-AM®	Orange oil extract	Physically compatible



Spruzit®	Pyrethrins + rapeseed oil	Observe a spray interval of 2 days before & after application
Toppel® 100	Cypermethrin	Physically compatible
Tracer®	Spinosad	Physically compatible
Verimark®	Cyantraniliprole	Physically compatible

ADVUVANTS

<u>BRAND NAME</u>	<u>ACTIVE INGREDIENT</u>	<u>COMMENTARY</u>
Activator® 90	Akylphenol ethoxylate, alcohol ethoxylate, tall oil fatty acids	Physically compatible
Breakthrough®	Polysiloxane	Physically compatible
Codacide	Oil seed rape extract	Observe a spray interval of 2 days before & after application
Silwet® L-77	Trisiloxane organosilicone copolymers	Physically compatible
Urea	Urea	Physically compatible 1%
Wetcit®	Alcohol Ethoxylate	Yes (0.8% v/v)

PLANT GROWTH REGULATORS

<u>BRAND NAME</u>	<u>ACTIVE INGREDIENT</u>	<u>COMMENTARY</u>
B-Nine®	Daminozide	Physically compatible
Bonzi®	Paclobutrazol	Physically compatible
Stabilan®	Chlormequat	Physically compatible

Please note if you wish to mix any herbicides with BotaniGard WP then please consult Certis UK.

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

For further information with regard to the warning phrases and symbols for this product please refer to the product label

Botanigard® is a registered trademark of Certis USA. Botanigard® contains Beauveria bassiana. Mapp no. 17054

Marketed by: Certis UK, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridge CB21 6AD

Tel: 0845 373 0305 Fax: 01223 891210

Scan here to download our free product guide app or visit www.certiseurope.co.uk from your smart phone or tablet

