

Best Use Practical Guidance - Potatoes

- Apply Crown MH from 3-5 weeks before haulm desiccation when more than 80% of tubers are larger than 25mm.
- Water Volume:
 - When dry, sunny and warm, use a minimum of 400L/ha of water in order to keep the crop as wet as possible.
 - Under normal moisture conditions and when during slow drying weather conditions (e.g. cloudy), a water volume of 250-300L/ha can be used.
- Crown MH is rainfast in 8-12 hours after application.
- Apply Crown MH as a solo application to allow optimum drying time and absorption.
- Recommend not to use with an adjuvant on potatoes, as adjuvants can speed up leaf drying time.
- DO NOT apply to stressed crops as uptake will be reduced.
 - If crop is water stressed, irrigate and wait 24 hours before applying.
- DO NOT apply when temperatures are above 25°C. If high daytime temperatures occur, spray in the evenings when temperature is cooling down.
- DO NOT apply to seed crops.



USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

For further information with regard to the warning phrases and symbols for this product please refer to the product label.

Crown[®] is a registered trademark of Kreglinger. Crown[®] contains maleic hydrazide. MAPP No. 18018.

Marketed by: Certis, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridge CB21 6AD

Tel : 0845 373 0305 Fax: 01223 891210 © Certis 2018