


Maximises Seed Health and Marketable Yield

- Effective on; dry rot (all strains of fusarium sp), silver scurf, gangrene, skin spot
- Excellent protectant. To obtain the best reduction of skin and wound diseases, apply before wounds begin to close.
- Maximum efficacy achieved when applied within 7-10 days after harvest
- Tubers should be clean and free from soil (less than 5% of surface)
- Treatment should be made before chitting to minimise mechanical damage



Programme Approach



For optimum efficacy of GAVEL:



Team Sprayers CTC 2 air treatment canopy

- Gavel is a protectant, therefore treat on store loading
- Apply to tubers passing over a roller table
- For best results use a canopy applicator with rotating arm nozzles (eg. TEAM CTC 2 air treatment canopy)
- Apply at 150ml/tonne in a total maximum solution of 2 litres water/tonne
- Treated seed must not be fed to livestock or go for human consumption

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

For further information with regard to the warning phrases and symbols for these products please refer to the product label.

Gavel® is a registered trademark of Certis Europe B.V. Gavel® contains imazalil. MAPP No. 17586.
Certis, 3 Riverside, Suite 5, Granta Park, Great Abington, Cambridge CB21 6AD. Tel: 0845 373 0305

www.certiseurope.co.uk Certis® 2019