

LATITUDE XL®
PCS 04721



SEED TREATMENT

A seed treatment for the reduction of take -all in winter and spring wheat, triticale and winter barley. This product is a flowable suspension concentrate containing 125 g/L (11.83% w/w) of silthiofam

Contents: XXX kg
Batch no:

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when bagging treated seed.

Environmental Protection

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/ Avoid contamination via drains from farmyards and roads)
To protect birds and wild mammals, treated seed should not be left on the soil surface. Bury or remove spillages

Treated seed should not be broadcast.

Seed should be drilled to a depth of 40mm into a well-prepared seed bed.

If seed is present on the soil surface, or if spills have occurred, then, if conditions are appropriate, the field should be harrowed and then rolled to ensure good incorporation.

Storage and disposal

Keep out of reach of children

Keep away from food, drink and animal feeding stuffs

Use appropriate containment to avoid environmental contamination

The material and its container must be disposed of in a safe way

LATITUDE XL®

A flowable suspension concentrate containing 125 g/Litre (11.83% w/w) of silthiofam



WARNING

MAY CAUSE DAMAGE TO ORGANS THROUGH PROLONGED OR REPEATED EXPOSURE
TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECT

Do not breathe/ dust/ fume/gas/ mist/ vapours/ sprays.

Avoid release to the environment.

Get medical advice/ attention if you feel unwell.

Collect spillage

Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty, clean containers which can be disposed of as non-hazardous waste

Contains methenamine 3-chloroallylochloride, 1,2-benzisothiazol-3(2H)-one, reaction mass of: 5-chloro-2-methyl-4-isothiazolin-3-one [EC no. 247-500-7] and 2-methyl-2H-isothiazol-3-one [EC no. 220-239-6] (3:1) May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 04721

IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE SEED TREATMENT

Crops: Winter and spring wheat, winter barley and triticale

Maximum individual dose: 200ml of product/100kg of seed

Maximum number of treatments: One per batch

Latest time of application: Immediately prior to drilling

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE: Refer to the label attached to the container
PROTECT FROM FROST

Marketed by:

CERTIS

Certis Europe BV, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire, CB21 6AD

Tel: 0044 (0)1223 894261 Fax: 0044 (0)1223 891210

E-mail: info@certiseurope.co.uk

For technical and non-emergency calls – phone 0044 (0)845 373 0305

For advice on medical emergencies, fires, spillages or chemical hazards only – phone 0870 190 6777

Approval holder: Certis Europe B.V, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire, CB21 6AD

®LATITUDE XL is a registered trademark of Mitsui & Co. Ltd.

CONDITIONS OF SUPPLY: The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumer's statutory rights are not affected.

SAFETY PRECAUTIONS FOR TREATED SEED BAG LABEL

THIS SEED HAS BEEN CHEMICALLY TREATED.

TREATED SEED must be drilled within the season of treatment

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when handling treated seed.

DO NOT HANDLE seed unnecessarily.

DO NOT USE TREATED SEED as food or feed.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE BEEN USED FOR TREATED SEED for food or feed.

KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.

WASH HANDS AND EXPOSED SKIN before meals and after work.

BURY OR REMOVE SPILLAGES.

HARMFUL TO GAME AND WILDLIFE. Treated seed should not be left on the soil surface, bury spillages

DO NOT SOW TREATED SEED FROM THE AIR.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions or Warnings

Seed quality: Do not use LATITUDE XL on seed with more than 16% moisture content, or on sprouted, cracked, or skinned seed. Test germination on all seed batches to ensure that the seed is viable and suitable for treatment with LATITUDE XL.

Disease controlled

LATITUDE XL contains silthiofam, a fungicide from the benzamide group. It is used in winter and spring wheat, winter barley and triticale as a fungicidal seed treatment for reducing the effects of take-all (*Gaeumannomyces graminis var tritici*)

LATITUDE XL can reduce both the severity of root rot and the incidence of whiteheads which can result in increased crop yield and grain quality. It is important to recognise that symptoms of take-all may be seen in treated crops in high disease situations.

Crop specific information

Rate of Use: 200ml product per 100kg of seed.

Application

In order to recirculate the product in the container, agitate the contents thoroughly before use.

Warning: containers of greater than 25 litres capacity should be handled only with mechanical assistance.

Apply LATITUDE XL simultaneously with a standard seed treatment using a recommended seed treatment machine which has been accurately calibrated to obtain the correct dose rate. Apply water, as necessary, to optimise coverage. Only apply LATITUDE XL using approved seed treatment equipment. For a list of approved equipment contact your Certis representative.

After use, thoroughly clean the application equipment using clean water.

Storage of Treated Seed

Drill seed treated with LATITUDE XL in the season of purchase. Although LATITUDE XL does not have any adverse effect on the storage life of treated seed, the viability of such seed and the performance of LATITUDE XL may be adversely affected by physical storage conditions. No liability can therefore be accepted for the performance of stored, treated seed.

Crops Grown for Processing

LATITUDE XL is unlikely to show any adverse effect on baking, malting, brewing and distilling processes. However, for crops grown for processing, consult your grain merchant or processor before use.

Drilling Instructions

Check drill calibration before use with each seed lot to ensure accurate seeding rates. Drill LATITUDE XL treated seed at a depth of 2.5-4 cm, into a well-prepared and firm seedbed, providing optimal conditions for crop emergence and establishment.

Use best practice to avoid adverse seedbed conditions such as very dry, fluffy, cloddy or capped seedbeds, water logged soil or too deep or too shallow drilling depths. Also avoid the use of seed with low germination capacity and/or vigour and residual herbicides that reduce field emergence.

Integrate the use of LATITUDE XL with cultural methods of control by optimising sowing date, seeding rate, nitrogen use and soil conditions (e.g. seed-bed preparation, drainage, pH etc).

Compatibility

LATITUDE XL shows excellent crop safety and compatibility with other seed treatments. It should be co-applied with standard seed treatments for control of seed-borne diseases.

For further information on the compatibility of Latitude XL with any other material consult your Certis representative.

Resistance Management

The possible development of resistance to LATITUDE XL cannot be excluded or predicted. Where such resistant disease strains may occur, LATITUDE XL is unlikely to give satisfactory control.

As a precaution do not treat more than three consecutive susceptible cereal crops in any one rotation. Also reduce resistance risk by adopting an integrated approach to take-all control

END OF LABEL TEXT

Material Safety Data Sheet

To access the Safety Data Sheet for this product, scan the QR code or use the web link below

URL: <https://cutt.ly/rvjN7v>



Alternatively, contact your supplier