

Magna<sup>®</sup> SL  
MAPP 19496



**A plant growth regulator, soluble concentrate formulation containing 270 g/L maleic hydrazide used to inhibit sprouting in onions and potatoes during storage and to control the growth of volunteer potatoes in succeeding crops.**

Batch no:

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS  
MAY APPLY TO THE USE OF THIS PRODUCT AT WORK

### **SAFETY PRECAUTIONS**

#### **Operator Protection**

WASH HANDS/ EXPOSED SKIN before eating and drinking/smoking/after work.

#### **Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

#### **Spray tank cleaning**

Wash application equipment thoroughly after use.

#### **Storage and disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into the spray tank and dispose of the container safely. Store product in a cool, dry place where it is not exposed to heat.

#### **Magna<sup>®</sup> SL**

A plant growth regulator, soluble concentrate formulation containing 270 g/L maleic hydrazide.

Dispose of content/ container to a licenced hazardous waste disposal contractor or collection site except for triple rinsed empty, clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions of use.**

**MAPP 19496**

<b>IMPORTANT INFORMATION</b>				
<b>FOR USE ONLY AS A PROFESSIONAL PLANT GROWTH REGULATOR</b>				
<b>Crops/ situations</b>	<b>Maximum individual dose</b>	<b>Maximum total dose</b>	<b>Maximum number of treatments</b>	<b>Latest time of application</b>
Bulb onions	8.9 L/ha	-	One per year	4 days before harvest
Potato	11 L/ha	-	One per year	21 days before harvest

Other specific restrictions:  
1) Do not apply to seed potato crops

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**



This label is compliant with the Voluntary Initiative Guidance

**Marketed by:** Certis

Certis Europe BV, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire, CB21 6AD

Tel: 0044 (0)1223 894261

Fax: 0044 (0)1223 891210

E-mail: [certis@certiseurope.co.uk](mailto:certis@certiseurope.co.uk)

**For technical and non-emergency calls – phone 0044 (0)845 373 0305**

**For advice on medical emergencies, fires, spillages or chemical hazards only – phone 0870 190 6777**

**Approval holder:** Kreglinger Europe NV, Grote Markt, 7, B-2000 Antwerp, Belgium,

®MAGNA is a registered trademark of Kreglinger

CONDITIONS OF SUPPLY: The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumer's statutory rights are not affected.

#### **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

**Purpose:**

MAGNA SL is a growth regulator which affects cell division, without affecting cell development. As a result, provided that the recommendations are followed, this product can be used for the purpose of inhibiting sprouting in onions and potatoes during storage. MAGNA SL helps inhibit growth of volunteer potatoes in succeeding crops.

**Preparing the spray solution:**

Fill the tank of spraying equipment half with water and turn on the agitation system. While the mixture is being agitated, add the required quantity of MAGNA SL to the tank and subsequently the rest of the required water. Ensure continuous agitation during spraying.

**Mixing with other products- general recommendations:**

Do not mix MAGNA SL with other growth regulators. When used on onions, absorption can be enhanced by adding a wetting agent.

**GENERAL RECOMMENDATIONS**

The product should only be applied to healthy plants (i.e. plants without diseases, pests, parasites or suffering drought stress)

The foliage of crops to be treated should be dry.

The best results will be obtained if spraying is conducted under cool conditions, not when the weather is dry and hot.

Do not use the product at temperatures above 26 °C. In the event of high temperatures, postpone application until cooler weather conditions occur, with temperatures below 26 °C and relative humidity of at least 50%.

Do not treat crops grown for seed

**BULB ONIONS**

Pre-harvest treatment will inhibit sprouting in store and enhance their quality.

**Application**

Apply using 500 to 600 litres of water per hectare for onions.

**Timing**

Apply MAGNA SL when the crop is approaching maturity, from 10% necking until no more than 50% leaf necking (BBCH 47-48) when the foliage is still green.

**Cautions**

'Spongy' bulbs may result if the spraying timing is not correct. Onions must be in good condition and properly cured if treatment is to be successful. All varieties of bulb onions can be treated. Salad and other types of onions should not be treated. Do not lift onions for at least four days after treatment. Do not spray earlier than 2 weeks before maturity. Do not apply to first year onion sets.

Do not use under drought conditions when the temperature exceeds 26°C or when the crop is under stress from other factors e.g. insect or disease damage. Avoid overlapping spray swaths.

**POTATOES**

Pre-harvest treatment to second early and main crop ware potatoes will inhibit sprouting in store and will help controlling growth of volunteer potatoes in succeeding crops.

**Application**

Apply using 300 to 600 litres of water per hectare for potatoes.

**Timing**

Apply from three to seven weeks before crop desiccation (Or topping/Pulverization) or the beginning of natural senescence (latest timing BBCH 89). The best result will be obtained from applications at the earlier timing. Apply MAGNA SL when sufficient tubers of size greater than 25 mm are present to give a marketable yield. Only tubers larger than 25 mm will continue to grow and result in a marketable yield. Some basal leaves may already be drying off at this stage.

**Cautions**

Occasional damage to terminal growth may occur following the application of MAGNA SL. Varieties with vigorous haulms and / or late maturing foliage may be prone to damage. All the second early and main crop varieties of potato can be treated. Tuber numbers and yield may be reduced if Magna SL is applied too early. Do not treat first early potatoes or potatoes being grown under polythene. Do not apply to potatoes within three weeks of haulm destruction or the onset of natural die back or that are suffering from pest, disease or climatic stress. Only treat healthy, actively growing crops. Do not apply to seed potato crops. MAGNA SL should be used on second-early and main crop varieties not being used for seed as part of an integrated programme for the control of volunteer potatoes in succeeding crops.

---

END OF LABEL TEXT

**Material Safety Data Sheet**

To access the Safety Data Sheet for this product, scan the QR code or use the web link below

URL: <https://cutt.ly/rvjN7v>



Alternatively, contact your supplier