



**RHINO® DSG  
MAPP 19730**

**GROUP    FRAC 7    FUNGICIDE**

**A powder for dry seed treatment formulation containing 60 g/kg flutolanil for the control of black scurf and some reduction of stem canker on seed potatoes.**

- Seed Potatoes

2 kg e

Batch no:

THE (COSHH) CONTROL OF SUBSTANCES HAZARDOUS TO HEALTH REGULATIONS MAY APPLY TO THE USE OF THIS PRODUCT AT WORK.

## **SAFETY PRECAUTIONS**

### **Operator protection**

This product must only be applied using automated powder dispensing equipment fitted on an automatic planter

Other engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the product.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, other engineering controls in addition to those specified above may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

TAKE OFF IMMEDIATELY all contaminated clothing.

DO NOT BREATHE DUST.

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating, drinking and smoking and after work.

### **Environmental protection**

DO NOT CONTAMINATE WATER WITH THE PRODUCT OR ITS CONTAINER. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads)

### **Storage and disposal**

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

BURY OR REMOVE SPILLAGES.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

Empty container completely and dispose of safely.

Sacks containing treated seed must not be used for food or feed'

**IMPORTANT INFORMATION**

**FOR USE ONLY AS AN AGRICULTURAL SEED TREATMENT**

Crop: Seed potatoes.  
Maximum individual dose: 2 kg of product per tonne of potatoes.  
Maximum number of treatments: One per batch  
Latest time of application: At planting

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**

**RHINO DSG**

A powder for dry seed treatment formulation containing 60 g/kg flutolanil



**WARNING**

**CAUSES SERIOUS EYE IRRITATION.  
HARMFUL TO AQUATIC LIFE WITH LONG LASTING EFFECTS.**

Wash hands thoroughly after handling.

Avoid release to the environment.

Wear protective gloves/protective clothing/eye protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

**MAPP 19730**

PROTECT FROM FROST.



This label is compliant with the CPA Voluntary Initiative Guidance

**Marketed by:**

**CERTIS**

Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridgeshire, CB21 6AD

Tel: 0044 (0)1223 894261

Fax: 0044 (0)1223 891210

E-mail: [certis@certiseurope.co.uk](mailto:certis@certiseurope.co.uk)

**For technical and non-emergency calls – phone 0044 (0)845 373 0305**

**For advice on medical emergencies, fires, spillages or chemical hazards ONLY  
– phone 0870 190 6777**

**Authorisation Holder:** Nichino Europe Co. Limited, 5 Pioneer Court, Vision Park,  
Histon, Cambridge, CB24 9PT

® Rhino is a registered trademark of Certis Europe BV

CONDITIONS OF SUPPLY: The Seller warrants that the goods shall at the time of delivery to the Buyer conform to the Seller's standard specification but all other conditions and warranties, whether express or implied by statute or custom of the trade or otherwise and whether as to condition, quality, performance, merchantability, fitness for any purpose or otherwise, are expressly excluded and, subject as aforesaid, the Seller shall be under no liability whatsoever, in contract or in tort, for or in respect of any loss or damage whatsoever resulting from or arising out of the goods or supply or use thereof, whether caused by the negligence of the Seller or otherwise. The Seller shall be under no liability in respect of the warranty given above unless the Buyer allows the Seller reasonable opportunity of inspecting the goods where practicable. A consumers statutory rights are not affected.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Rhino DSG controls black scurf and some reduction of stem canker on potatoes, when applied to the seed at planting

## **Crops**

Rhino DSG can be used on seed potatoes for the seed, ware or processing market. It can be used on the seed tubers of all varieties of early, second early and maincrop

potatoes grown for ware or processing. Check with your processor before using Rhino DSG on the crop.

## Timing

Rhino DSG should be applied to the seed tubers at planting.

## Disease Control

Rhino DSG is a seed potato treatment which controls black scurf (*Rhizoctonia solani*) and also provides some reduction of stem canker. Where soil borne inoculum is high, treatment of seed tubers with Rhino DSG may not give full protection against infection of stems and daughter tubers.

## Application

### DOSE

Use Rhino DSG at 2 kg per tonne of seed potatoes.

### CROP CONDITIONS

For best results the tubers should be clean and free from soil (less than 5% of surface). Disease under areas of soil/dust will not be exposed to dust and will not be controlled. Seed tubers should be of good quality and vigour (free from bacterial rots, physical damage, virus infection etc), and should not be sprouted to such an extent that mechanical damage to the shoots will occur during treatment or planting.

### APPLICATION TECHNIQUES

**On-planter application** - specialised applicators (e.g. Team Sprayers applicator) must be attached to planters to distribute the required amount of product onto tubers as they are being planted.

## Storage

Store above 5° C

---

### Section 6 of the Health and Safety at Work Act

#### Additional Product Safety Information

(This section does not form part of the product label under the Control of Pesticide Regulations 1986).

The product label provides information on a specific pesticidal use of the product; do not use otherwise, unless you have assessed any potential hazard involved, the safety measures required and that particular use has "off-label" approval or is otherwise permitted under the Control of Pesticide Regulations. The information on this label is based on the best available information including data from test results.

### **Material Safety Data Sheet**

To access the Safety Data Sheet for this product, scan the QR code or use the web link below

URL: <https://cutt.ly/rvjN7v>



Alternatively, contact your supplier

### **CASE LABEL TEXT**

---

**RHINO DSG**

8 x 2 kg

1109TEXT