

## Novel fungicide for powdery mildew control EAMU for use on strawberries

- New active ingredient and mode of action for strawberry crops
- Preventative or curative activity depending on pathogen pressure and concentration used
- Translaminar and vapour movement for longer protection and better coverage of treated leaves
- Short harvest interval (three days)
- Wide water volume range. 600 - 1500 L
- Two opportunities to target mildew


### Using Takumi SC

- Use in programme with other mildew products.
- Use high concentration on young plants when disease is first seen.
- Mature plants may require higher water volumes than specified on the label, providing the label rate of 150ml/Ha is not exceeded.
- Use as part of a robust anti-resistance strategy and combine Takumi SC with products containing varying modes of action such as Karma.

### Beneficial insect safety

- Takumi SC has low impact on beneficial insects such as *Encarsia formosa* and *Macrolophus caliginosus*.
- Takumi SC can easily be integrated into an IPM programme.
- Contact Certis Europe for further details

Powdery mildew control in strawberries compared to untreated %



<b>Formulation</b>	A suspension concentrate (SC) containing 100 g/L cyflufenamid		
<b>Chemical class</b>	Benzamidoxime	<b>FRAC</b>	U6
<b>Crops</b>	Strawberries		
<b>No. applications</b>	2 (per crop), avoid consecutive applications. Follow FRAC guidelines. Use in a programme with other products with different mode of action.		
<b>Max dose rate</b>	0.15 L/Ha	<b>Harvest Period</b>	3 Days
<b>Water volume</b>	Preventative treatment up to 1500 L/Ha, curative treatment 600 L/Ha	<b>MAPP No.</b>	16000
		<b>EMAU No.</b>	2055 of 2016



USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE

For further information with regard to the warning phrases and symbols for these products please refer to the product labels.  
 Extension of Use: This extension of the authorised use provides for the use of TAKUMI SC<sup>®</sup> (M16000) in respect of crops and situations, other than those included on the product label. No efficacy or phytotoxicity data have been assessed and as such the 'extension of use', is at all times done at the user's choosing, and the commercial risk is entirely theirs. Users must be in possession of a copy of the Extension of Authorisation Number: 2055 of 2016 for full details of product extension prior to use.  
 TAKUMI SC<sup>®</sup> is a registered trademark of Nippon Soda Co., Ltd. Marketed by: Certis, Suite 5, 3 Riverside, Granta Park, Great Abington, Tel: 0845 373 0305 • Email: [infocertisuk@certiseurope.com](mailto:infocertisuk@certiseurope.com) • [www.certiseurope.co.uk](http://www.certiseurope.co.uk)  
 08/2017

[scan here to download our free product guide app](#)

or visit [m.certiseurope.co.uk](http://m.certiseurope.co.uk) from your smart phone or tablet