



TOPSIN SCLEROTINIA UPDATE: April 2018

- Although not seen in crops every year, Sclerotinia can lead to yield losses of up to 60% when conditions favour the disease.
- The AHDB Sclerotinia Risk Alerts for 2018 are already showing that sites across the west, south and east are at high risk of disease development.
- When risk is high and the flowering period is prolonged, a two spray programme or robust mid-flowering spray, may be appropriate.
- TOPSIN (containing thiophanate-methyl) can be a useful tool for the agronomist and grower, giving strong cost-effective Sclerotinia control alone, or in tank-mix with a product with a different mode of action.
- TOPSIN can be used as part of a 2 spray programme alone at up to 0.71kg/Ha or in tank-mix at 0.5kg/Ha.
- Alternatively, if only one spray is planned mid-flower, TOPSIN can be a very useful addition, giving good efficacy and helping with resistance management.
- For more information on use and tank-mixes, contact your local Certis advisor or call our technical helpline on 0845 373 0305.
- For more information about the Sclerotinia risk in your area, please visit:

<https://cereals.ahdb.org.uk/monitoring/sclerotinia/sclerotinia-risk-report.aspx>

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

For further information with regard to the warning phrases and symbols for this product please refer to the product label.
Topsin® is a registered trademark of Nippon Soda Co. Ltd., Japan. Topsin® WG contains thiophanate-methyl. MAPP no. 13988
Marketed by: Certis, Suite 5, 3 Riverside, Granta Park, Great Abington, Cambridge CB21 6AD. © Certis 2016
Tel : 0845 373 0305 • Fax: 01223 891210

CERTIS

INNOVATIVE SOLUTIONS FOR CROP PROTECTION

Tel: 0845 373 0305 • Email: infocertisuk@certiseurope.com

www.certiseurope.co.uk

